

SUPPLEMENTARY MATERIALS

Literature search:

Search criteria were as follow:

- pain[Majr] AND chronic AND (tramadol OR tapentadol) AND placebo
- tramadol OR tapentadol AND ("Pain Measurement" OR "Pain"[Majr]) NOT neoplasms AND placebo AND drug therapy AND Humans
- tramadol OR tapentadol AND pain AND (publisher OR "in process") AND trial

These searches resulted in a list of sources of potential interest. This list was incorporated into Reference Manager® Version 11 (Thomson Reuters, Carlsbad, CA). Further review reduced the list to sources of primary interest. Each of these sources was individually examined to identify or rule out potential additional sources of information.

The sources of primary interest identified above were used to create a database in an Excel® spreadsheet format. Studies were included in the database if they reflected treatments, study conditions, and patient populations of interest, and were rejected if they contained information not of interest or if the reports were insufficiently detailed to be used to construct a model. In some cases, a study contained data of interest and additional data not deemed to be of interest. In these cases only the data of interest were extracted. The following inclusion criteria were used regarding studies to be entered into the database:

- Active treatment with tramadol or tapentadol
- Preferences for double-blind, randomized, controlled, parallel group study design, but may include cross-over and open-label design in some cases if other criteria are met
- Contained data reflective of pain, pain intensity, patient severity, patient's global assessment, physician's global assessment, adverse events (gastrointestinal) and dropouts.

Study results from tables, figures, or textual information in the individual reports were included. Relevant figures were digitized, and then converted into numeric values using Didger® Version 3 (Golden Software, Golden, CO). Information pertaining to the different study design settings and the patient characteristics were captured in the database to the extent possible.

In all cases where available relevant summary statistics (mean, median, geometric mean, harmonic mean) and the corresponding measurements of standard deviation (SD) or standard error (SE) were available, they were entered into the database. In some cases SD/SE values were either not provided or not able to be digitized with sufficient accuracy due to poor source document resolution

Included studies were randomized, placebo-controlled, with a parallel-group or cross-over design, published in 1985 or later, with at least 5 days of treatment exposure or follow-up that reported pain intensity.

Table S1: List of trials retained in the analysis, with key information

Author	Pub Year	Duration	N*	Pain intensity	Constipation	Nausea	Dizziness	Vomiting	Somnolence	DropOut for AE	DropOut for LoE	Dosing Method	Treatment arms retained in the current analysis
Adler	2002	4 weeks	202	✓						✓	✓	titration	tramadol
Affilalo	2010	15 weeks	681	✓	✓	✓	✓	✓	✓	✓	✓	titration to fixed	tapentadol / placebo
Babul	2004	12 weeks	246	✓	✓	✓	✓	✓	✓	✓	✓	titration	tramadol / placebo
Beaulieu	2007	4 weeks	122		✓	✓	✓	✓	✓	✓	✓	flexible	tramadol
Bennett	2003	91 days	313	✓	✓	✓	✓		✓	✓	✓	flexible	tramadol / placebo
Bianchi	2003	1 week	8	✓						✓		fixed	tramadol / placebo
Biovail Laboratories Inc	2005	12 weeks	246	✓						✓	✓	titration to fixed	tramadol
Bureau	2003	6 weeks	127	✓								flexible	tramadol / placebo
Burch	2007	12 weeks	589	✓	✓	✓	✓		✓	✓	✓	titration to fixed	tramadol / placebo
Buynak	2010	15 weeks	637	✓	✓	✓	✓	✓	✓	✓	✓	titration to fixed	tapentadol / placebo
Caldwell	1999	30 days	36	✓	✓	✓	✓	✓	✓	✓	✓	titration	placebo
Chindalore	2005	3 weeks	51	✓	✓	✓	✓	✓	✓	✓	✓	escalation	placebo
Choi	2007	2 weeks	189	✓	✓	✓	✓	✓	✓	✓		titration	tramadol
Dalgin	1997	39 days	146	✓	✓	✓						titration	tramadol
Elan	2002	4 weeks	67	✓								fixed	placebo
Emkey	2004	91 days	306	✓	✓	✓	✓	✓	✓	✓	✓	titration	tramadol / placebo
Fishman	2007	12 weeks	552		✓	✓	✓	✓	✓	✓	✓	titration to fixed	tramadol / placebo
Fleischmann	2001	91 days	129	✓	✓	✓	✓			✓	✓	titration	tramadol / placebo
Freeman	2007	9 weeks	312	✓	✓	✓	✓		✓	✓	✓	flexible	tramadol / placebo
Gana	2006	12 weeks	1011	✓	✓	✓	✓	✓	✓	✓	✓	titration to fixed	tramadol / placebo
Gimbel	2003	6 weeks	77	✓	✓	✓	✓	✓	✓	✓	✓	flexible	placebo
Hale	2005	18 days	75	✓	✓					✓	✓	flexible	placebo
Hale	2009	12 weeks	679	✓	✓	✓	✓	✓	✓	✓		flexible	tapentadol
Harati	1998	42 days	131	✓	✓	✓	✓	✓	✓	✓	✓	flexible	tramadol / placebo

Hartrick	2009	6 weeks	488		✓	✓	✓	✓	✓	✓	flexible	tapentadol / placebo
Jensen	1994	2 weeks	81		✓	✓	✓	✓		✓	fixed	tramadol
Kelly	2009	15 weeks	681	✓	✓	✓	✓	✓	✓		flexible	tapentadol / placebo
Kelly	2010	15 weeks	344							✓	flexible	tapentadol
Kosinski	2007	12 weeks	1011	✓							fixed	tramadol / placebo
Lee	2006	1 week	267	✓						✓	fixed	tramadol / placebo
Li	2008	7 days	48	✓	✓	✓		✓		✓	fixed	tramadol
Ma	2008	4 weeks	58	✓	✓	✓	✓	✓	✓		flexible	placebo
Malonne	2004	14 days	230	✓	✓	✓	✓	✓	✓	✓	fixed	tramadol / placebo
Markenson	2005	90 days	51	✓	✓	✓	✓	✓	✓	✓	flexible	placebo
Matsumoto	2005	4 weeks	124		✓	✓	✓	✓	✓	✓	fixed	placebo
Mongin	2004	12 weeks	314		✓	✓		✓	✓	✓	titration	tramadol
Mullican	2001	4 weeks	309		✓	✓	✓	✓	✓	✓	titration	tramadol
Norrbrink	2009	4 weeks	35		✓	✓	✓				flexible	tramadol / placebo
Pavelka	1998	4 weeks	54							✓	prn use	tramadol
Peloso	2000	4 weeks	35	✓							fixed	placebo
Peloso	2004	91days	336	✓	✓	✓	✓	✓	✓	✓	titration	tramadol / placebo
Rauck	1994	4 weeks	234	✓	✓	✓	✓	✓	✓	✓	titration	tramadol
Roth	2000	18 months	45	✓	✓	✓	✓	✓	✓	✓	fixed	placebo
Ruoff	2003	91 days	318	✓						✓	titration	tramadol / placebo
Schnitzer	2000	7 weeks	254			✓				✓	flexible	tramadol / placebo
Schwartz	2011	12 weeks	389	✓	✓	✓	✓	✓		✓	fixed	tapentadol / placebo
Silverfield	2002	10 days	212	✓	✓	✓	✓	✓	✓	✓	flexible	tramadol / placebo
Sindrup	1999	4 weeks	90		✓	✓	✓				flexible	tramadol / placebo
Thorne	2008	8 weeks	154	✓	✓	✓	✓	✓	✓	✓	titration	tramadol / placebo
Vidal	2005	6 weeks	132							✓	titration to fixed	tramadol
Vorsanger	2008	12 weeks	384		✓	✓	✓			✓	fixed	tramadol / placebo
Webster	2006	12 weeks	101	✓	✓	✓	✓	✓	✓	✓	flexible	placebo
Wild	2010	52 weeks	894	✓	✓	✓	✓	✓	✓	✓	flexible	tapentadol
Contributing articles				39	40	40	36	31	31	44	36	

N*: maximum between number of patients contributing to the efficacy (PI) analysis or safety analysis (AEs and DropOuts)

Table S2: Consistency between previous and current meta-analysis results concerning frequency of withdrawals and of specific adverse events

	Moore and McQuay (2005) – Table 3		Current analysis		
	placebo	oral opioids	placebo	tapentadol	tramadol
Withdrawals					
Due to adverse event	7.1	22	7.1	18.7	20.5
Due to lack of efficacy	20	6.5	18.5	6.1	7.8
Specific adverse events					
Constipation	5.0	15	5.3	15.1	18.0
Nausea	5.6	21	8.0	21.7	22.2
Vomiting	2.4	10	2.2	6.7	9.9
Somnolence	4.0	14	3.8	12.6	13.2
Dizziness	4.5	14	4.6	15.7	13.2

Table S3: Reference list

1. Adler L., McDonald C., O'Brien C., *et al.* A comparison of once-daily tramadol with normal release tramadol in the treatment of pain in osteoarthritis. *Journal of Rheumatology*, 2002, **29**: 2196-2199
2. Afilalo M., Etropolski M.S., Kuperwasser B., *et al.* Efficacy and safety of tapentadol extended release compared with oxycodone controlled release for the management of moderate to severe chronic pain related to osteoarthritis of the knee. *Clinical Drug Investigation*, 2010, **30**: 489-505
3. Babul N., Noveck R., Chipman H., *et al.* Efficacy and safety of extended-release, once-daily tramadol in chronic pain: a randomized 12-week clinical trial in osteoarthritis of the knee. *Journal of Pain and Symptom Management*, 2004, **28**: 59-71
4. Beaulieu A., Peloso P., Bensen W., *et al.* A randomized, double-blind, 8-week crossover study of once-daily controlled-release tramadol versus immediate-release tramadol taken as needed for chronic noncancer pain. *Clinical Therapeutics*, 2007, **29**: 49-60
5. Bennett R.M., Kamin M., Karim R., *et al.* Tramadol and acetaminophen combination tablets in the treatment of fibromyalgia pain: a double-blind, randomized, placebo-controlled study. *American Journal of Medicine*, 2003, **114**: 537-545
6. Bianchi M., Brogginini M., Balzarini P., *et al.* Effects of tramadol on synovial fluid concentration of substance P and interleukin-6 in patients with knee osteoarthritis: comparison with paracetamol. *International Immuno-pharmacology*, 2003, **3**: 1901-1908
7. Boureau F., Legallicier P., Kabir-Ahmadi M. Tramadol in post-herpetic neuralgia: a randomized, double-blind, placebo-controlled trial. *Pain*, 2003, **104**: 323-331.
8. Burch F., Fishman R., Messina N., *et al.* A comparison of the analgesic efficacy of tramadol contramid OAD versus placebo in patients with pain due to osteoarthritis. *Journal of Pain and Symptom Management*, 2007, **34**: 328-338
9. Buyank R., Shapiro D.Y., Okamoto A., *et al.* Efficacy and safety of tapentadol extended release for the management of chronic low back pain: results of a prospective, randomized, double-blind, placebo-and active-controlled phase III study. *Expert Opinions in Pharmacotherapy*, 2010, **11**: 1787-1804
10. Caldwell J.R., Hale M.E., Boyd R.E., *et al.* Treatment of osteoarthritis pain with controlled release oxycodone or fixed combination oxycodone plus acetaminophen added to non steroidal anti-inflammatory drugs: a double blind, randomized, multicenter, placebo controlled trial. *Journal of Rheumatology*, 1999, **26**: 862-869
11. Chindalore V.L., Craven R.A., Yu K.P., *et al.* Adding ultralow-dose naltrexone to oxycodone enhances and prolongs analgesia: a randomized, controlled trial of Oxytrex. *The Journal of Pain*, 2005, **6**: 392-399
12. Choi C.-B., Song J.S., Kang Y.M., *et al.* A 2-week, multicenter, randomized, double-blind, double-dummy, add-on study of the effects of titration on tolerability of tramadol/acetaminophen combination tablet in Korean adults with knee osteoarthritis pain. *Clinical Therapeutics*, 2007, **29**: 1381-1389
13. Dalgin P., Comparison of tramadol and ibuprofen for the chronic pain of osteoarthritis. *Arthritis Rheum.*, 1997, **40**: S86

14. Emkey R., Rosenthal N., Wu S.-C., *et al.* Efficacy and safety of tramadol/acetaminophen tablets (Ultracet®) as add-on therapy for osteoarthritis pain in subjects receiving a COX-2 nonsteroidal anti-inflammatory drug: a multicenter, randomized, double-blind, placebo-controlled trial. *Journal of Rheumatology*, 2004, **31**: 150-156
15. FDA CDER. Clinical review of Avinza® Morphine sulfate extended release capsules for the management of patients requiring continuous opioid analgesia – Sponsor: Elan Pharmaceutical / Ligand Pharmaceuticals. NDA #21-260, Medical review, 2002, March 20.
http://www.accessdata.fda.gov/drugsatfda_docs/nda/2002/21-260_Avinza_Medr_P1.pdf
16. FDA CDER. Clinical review of Raliva® Tramadol hydrochloride extended release tablets (100, 200 and 300 mg) for the management of moderate to moderately severe chronic pain in adults – Sponsor: Biovail Laboratories Inc. NDA #21-692, Medical review, 2005, August 23.
http://www.accessdata.fda.gov/drugsatfda_docs/nda/2005/021692s000_Raliva_MedR.pdf
17. Fishman R.L., Kistler C.J., Ellerbusch M.T., *et al.* Efficacy and safety of 12 weeks of osteoarthritic pain therapy once-daily tramadol (Tramadol Contramid OAD). *Journal of Opioid Management*, 2007, **3**: 273-280
18. Fleischmann R.M., Caldwell J.R., Roth S.H., *et al.* Tramadol for the treatment of joint pain associated with osteoarthritis: a randomized, double-blind, placebo-controlled trial. *Current Therapeutic Research*, 2001, **62**: 113-128
19. Freeman R., Raskin P., Hewitt D.J., *et al.* Randomized study of tramadol/acetaminophen versus placebo in painful diabetic peripheral neuropathy. *Current Medical Research and Opinion*, 2007, **23**: 147-161
20. Gana T.J., Pascual M.L.G., Fleming R.R.B., *et al.* Extended-release tramadol in the treatment of osteoarthritis: a multicenter, randomized, double-blind, placebo-controlled, clinical trial. *Current Medical Research and Opinion*, 2006, **22**: 1391-1401
21. Gimbel J.S., Richards P., Portenoy R.K., Controlled-release oxycodone for pain in diabetic neuropathy: a randomized controlled trial. *Neurology*, 2003, **60**: 927-934
22. Hale M., Dvergsten C., Gimbel J., Efficacy and safety of oxymorphone extended release in chronic low back pain: Results of a randomized, double-blind, placebo- and active-controlled phase III study. *The Journal of Pain*, 2005, **6**: 21-28
23. Hale M., Upmalis D., Okamoto A., *et al.* Tolerability of tapentadol immediate release in patients with lower back pain or osteoarthritis of the hip or knee over 90 days: a randomized, double-blind study. *Current Medical Research and Opinion*, 2009, **25**: 1095-1104
24. Harati Y., Gooch C., Swenson M., *et al.* Double-blind randomized trial of tramadol for the treatment of the pain of diabetic neuropathy. *Neurology*, 1998, **50**: 1842-1846
25. Hartrick C., Van Hove I., Stegmann J.-U., *et al.* Efficacy and tolerability of tapentadol immediate release and oxycodone HCl immediate release in patients awaiting primary joint replacement surgery for end-stage joint disease: a 10 day, phase III, randomized, double-blind, active- and placebo controlled study. *Clinical Therapeutics*, 2009, **31**: 260-271

26. Jensen E.M., Ginsberg F., Tramadol versus dextropropoxyphene in the treatment of osteoarthritis - a short term double-blind study. *Drug Investigation*, 1994, **8**: 211-218
27. Kelly K., Kuperwasser B., Okamoto A., *et al.* Efficacy and gastrointestinal tolerability of tapentadol extended release in a randomized, double-blind, placebo-and active-controlled study in patients with moderate to severe chronic osteoarthritis knee pain. *5th World Congress of the World Institute of Pain, March 13-16, New York, USA, 2009.*
28. Kelly K., Lange B., Etropolski M.S., *et al.* Dose stability of tapentadol extended release (ER) for the relief of moderate-to-severe chronic osteoarthritis knee pain. *AAPM annual meeting 2010*, 2010, abstract 125.
29. Kosinski M., Janagap C., Gajria K., *et al.* Pain relief and pain-related sleep disturbance with extended-release tramadol in patients with osteoarthritis. *Current Medical Research and Opinion*, 2007, **23**: 1615-1626
30. Lee E.Y., Lee E.B., Park B.J., *et al.* Tramadol 37.5-mg/acetaminophen 325-mg combination tablets added to regular therapy for rheumatoid arthritis pain: a 1-week, randomized, double-blind, placebo-controlled trial. *Clinical Therapeutics*, 2006, **28**: 2052-2060
31. Li C., Ni J., Wang Z., *et al.* Analgesic efficacy and tolerability of flupirtine vs. tramadol in patients with sub-acute low back pain: a double-blind multi-centre trial. *Current Medical Research and Opinion*, 2008, **24**: 3523-3530
32. Ma K., Jiang W., Zhou Q., *et al.* The efficacy of oxycodone for management of acute pain episodes in chronic neck pain patients. *International Journal of Clinical Practice*, 2008, **62**: 241-247
33. Malonne H., Coffiner M., Sonet B., *et al.* Efficacy and tolerability of sustained-release tramadol in the treatment of symptomatic osteoarthritis of the hip or knee: a multicenter, randomized, double-blind, placebo-controlled study. *Clinical Therapeutics*, 2004, **26**: 1774-1782
34. Markenson J.A., Croft J., Zhang P.G., *et al.* Treatment of persistent pain associated with osteoarthritis with controlled-release oxycodone tablets in a randomized controlled clinical trial. *The Clinical Journal of Pain*, 2005, **21**: 524-535
35. Matsumoto A., Babul N., Ahdieh H., Oxymorphone extended-release tablets relieve moderate to severe pain and improve physical function in osteoarthritis: results of a randomized, double-blind, placebo- and active-controlled phase III trial. *Pain Medicine*, 2005, **6**: 357-366
36. Mongin G., Yakusevich V., Köpe N., *et al.* Efficacy and safety assessment of a novel once-daily tablet formulation of tramadol. *Clinical Drug Investigation*, 2004, **24**: 545-558
37. Mullican W.S., Lacy J.R., Tramadol/acetaminophen combination tablets and codeine/acetaminophen combination capsules for the management of chronic pain: a comparative trial. *Clinical Therapeutics*, 2001, **23**: 1429-1445
38. Norrbrink C., Lundeberg T., Tramadol in neuropathic pain after spinal cord injury: a randomized, double-blind, placebo-controlled trial. *The Clinical Journal of Pain*, 2009, **25**: 177-184
39. Pavelka K., Peliskova Z., Stehlikova H., *et al.* Intraindividual differences in pain relief and functional improvement in osteoarthritis with diclofenac or tramadol. *Clinical Drug Investigation*, 1998, **16**: 421-429

40. Peloso P.M., Bellamy N., Bensen W., et al. Double blind randomized placebo control trial of controlled release codeine in the treatment of osteoarthritis of the hip or knee. *Journal of Rheumatology*, 2000, **27**: 764-771
41. Peloso P.M., Fortin L., Beaulieu A., et al. Analgesic efficacy and safety of tramadol/acetaminophen combination tablets (Ultracet®) in treatment of chronic low back pain: a multicenter, outpatient, randomized, double-blind, placebo, controlled trial. *Journal of Rheumatology*, 2004, **31**: 2454-2463
42. Rauck R.L., Ruoff G.E., McMillen J.I., Comparison of tramadol and acetaminophen with codeine for long-term pain management in elderly patients. *Current Therapeutic Research*, 1994, **55**: 1417-1431
43. Roth S.H., Fleischmann R.M., Burch F.X., et al. Around-the-clock, controlled-release oxycodone therapy for osteoarthritis-related pain: placebo-controlled trial and long-term evaluation. *Archives of Internal Medicine*, 2000, **160**: 853-860
44. Ruoff G.E., Rosenthal N., Jordan D., et al. Tramadol/acetaminophen combination tablets for the treatment of chronic lower back pain: a multicenter, randomized, double-blind, placebo-controlled outpatient study. *Clinical Therapeutics*, 2003, **25**: 1123-1141
45. Schnitzer T.J., Gray W.L., Paster R.Z., et al. Efficacy of tramadol in treatment of chronic low back pain. *Journal of Rheumatology*, 2000, **27**: 772-778
46. Schwartz S., Etropolski M.S., Shapiro D.Y., et al. Safety and efficacy of tapentadol ER in patients with painful diabetic peripheral neuropathy: results of a randomized-withdrawal, placebo-controlled trial. *Current Medical Research and Opinion*, 2011, **27**: 151-162
47. Silverfield J.C., Kamin M., Wu S.-C., et al. Tramadol/acetaminophen combination tablets for the treatment of osteoarthritis flare pain: a multicenter, outpatient, randomized, double-blind, placebo-controlled, parallel-group, add-on study. *Clinical Therapeutics*, 2002, **24**: 282-297
48. Sindrup S., Andersen G., Madsen C., et al. Tramadol relieves pain and allodynia in polyneuropathy: a randomised, double-blind, controlled trial. *Pain*, 1999, **83**: 85-90
49. Thorne C., Beaulieu A.D., Callaghan D.J., et al. A randomized, double-blind, crossover comparison of the efficacy and safety of oral controlled release tramadol and placebo in patients with painful osteoarthritis. *Pain Research and Management*, 2008, **13**: 93-102
50. Vidal J., Carro A., Ponce A., et al. Efficacy and safety of tramadol alone vs tramadol plus naproxen or naproxen alone, for the treatment of chronic pain associated to hip or knee-joint osteoarthritis non-responsive to optimum and stable doses of NSAIDS. *Arthritis and Rheumatism*, 2005, **52**: S508
51. Vorsanger G.J., Xiang J., Gana T.J., et al. Extended-release tramadol (tramadol ER) in the treatment of chronic low back pain. *Journal of Opioid Management*, 2008, **4**: 87-97
52. Webster L.R., Butera P.G., Moran L.V., et al. Oxytrex minimizes physical dependence while providing effective analgesia: a randomized controlled trial in low back pain. *The Journal of Pain*, 2006, **7**: 937-946
53. Wild J.E., Grond S., Kuperwasser B., et al. Long-term safety and tolerability of tapentadol extended release for the management of chronic low back pain or osteoarthritis pain. *Pain Practice*, 2010, **10**: 416-427